arXiv:quant-ph/0301145v2 30 Jan 2003

Two–Level System and Some Approximate Solutions in the Strong Coupling Regime

Kazuyuki FUJII *
Department of Mathematical Sciences
Yokohama City University
Yokohama, 236-0027
Japan

Abstract

In this paper we treat the 2–level system interacting with external fields without the rotating wave approximation and construct some approximate solutions in the strong coupling regime.

In this paper we consider the 2–level system interacting with external fields (a radiation field for example) and treat this system without the rotating wave approximation (RWA). Usually we assume the RWA and solve the equation. This approximation is appropriate in the weak coupling regime, however in the strong coupling one it is not applicable. See for example [1] and [2]. How can we treat the system in the strong coupling regime ? Shahriar et al treated this problem in [3], [4] and [5] and constructed some approximate solutions which contained what they called the Bloch–Siegert oscillation. See also [6], [7], [8], [9] as another approachs.

---

*E-mail address : fujii@yokohama-cu.ac.jp

By the way we have treated some important models coming from quantum optics in the strong coupling regime, see [10], [12] and [13] (also [15] for an experiment in the strong coupling regime). See [11] as a general introduction in this field. This method is also applicable in the case, so we construct some approximate solutions which are completely different from those of Shahriar et al. We believe that our solutions are more general.

Let $\{\sigma_1, \sigma_2, \sigma_3\}$ be Pauli matrices :

$$\sigma_1 = \left(\begin{array}{cc} 0 & 1 \\ 1 & 0 \end{array}\right), \quad \sigma_2 = \left(\begin{array}{cc} 0 & -i \\ i & 0 \end{array}\right), \quad \sigma_3 = \left(\begin{array}{cc} 1 & 0 \\ 0 & -1 \end{array}\right), \tag{1}$$

and $\sigma_+ = (1/2)(\sigma_1 + i\sigma_1)$, $\sigma_- = (1/2)(\sigma_1 - i\sigma_1)$.

We consider an atom with two energy levels which interacts with external field with $g\cos(\omega t)$. The Hamiltonian in the dipole approximation is given by

$$H = -\frac{\Delta}{2}\sigma_3 + g\ \cos(\omega t)\sigma_1, \quad \cos(\omega t) = (1/2)(\mathrm{e}^{i\omega t} + \mathrm{e}^{-i\omega t}), \tag{2}$$

where ω is the frequency of the external field, Δ the energy difference between two levels of the atom, g the coupling constant between the external field and the atom. In the following we cannot assume the rotating wave approximation (which neglects the fast oscillating terms), namely the Hamiltonian given by

$$H = -\frac{\Delta}{2}\sigma_3 + \frac{g}{2}\left(\mathrm{e}^{i\omega t}\sigma_+ + \mathrm{e}^{-i\omega t}\sigma_-\right). \tag{3}$$

Mysteriously enough we cannot solve this simple model completely (maybe non–integrable), nevertheless we have found this model has a very rich structure.

We would like to solve the Schrödinger equation with the Hamiltonian (2) in **the strong coupling regime** $g \gg \Delta$. Therefore we change in (2) a role of the kinetic term and the interaction term like

$$H = H_0 - \frac{\Delta}{2}\sigma_3 \equiv g\ \cos(\omega t)\sigma_1 - \frac{\Delta}{2}\sigma_3.$$

First we solve the interaction term which is an easy task. Before that let us make some preliminaries.

Let W be a Walsh–Hadamard matrix

$$W = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} = W^{-1} \tag{4}$$

then we can diagonalize σ_1 by using this W as $\sigma_1 = W\sigma_3 W^{-1}$. The eigenvalues of σ_1 is $\{1, -1\}$ with eigenvectors

$$|1\rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \quad |-1\rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ -1 \end{pmatrix}.$$

We note that $\sigma_3|1\rangle = |-1\rangle$, $\sigma_3|-1\rangle = |1\rangle$ and

$$|1\rangle\langle 1| = \frac{1}{2} \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} = W \begin{pmatrix} 1 & \\ & 0 \end{pmatrix} W = W(|0)(0|)W,$$

$$|-1\rangle\langle -1| = \frac{1}{2} \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix} = W \begin{pmatrix} 0 & \\ & 1 \end{pmatrix} W = W(|1)(1|)W,$$

where $|0) = (1, 0)^{\mathrm{t}}$ and $|1) = (0, 1)^{\mathrm{t}}$.

Then it is easy to see

$$H_0 = g\cos(\omega t)|1\rangle\langle 1| - g\cos(\omega t)|-1\rangle\langle -1|, \tag{5}$$

so the Schrödinger equation $i\frac{d}{dt}\psi = H_0\psi$ can be solved to be

$$\psi(t) = \left\{ \mathrm{e}^{-i\frac{g}{\omega}sin(\omega t)}|1\rangle\langle 1| + \mathrm{e}^{i\frac{g}{\omega}sin(\omega t)}|-1\rangle\langle -1| \right\} \psi_0, \tag{6}$$

where ψ_0 is a constant vector. Since we want to solve the full Schrödinger equation

$$i\frac{d}{dt}\psi = \left(H_0 - \frac{\Delta}{2}\sigma_3 \right) \psi, \tag{7}$$

we appeal to the **method of constant variation**. Namely under $\psi_0 \longrightarrow \psi_0(t)$ in (6) we substitute (6) into (7), then we obtain

$$\begin{aligned} i\frac{d}{dt}\psi_0 = -\frac{\Delta}{2} \times & \\ \left\{ \mathrm{e}^{i\frac{g}{\omega}sin(\omega t)}|1\rangle\langle 1| + \mathrm{e}^{-i\frac{g}{\omega}sin(\omega t)}|-1\rangle\langle -1| \right\} & \sigma_3 \left\{ \mathrm{e}^{-i\frac{g}{\omega}sin(\omega t)}|1\rangle\langle 1| + \mathrm{e}^{i\frac{g}{\omega}sin(\omega t)}|-1\rangle\langle -1| \right\} \psi_0 \\ = -\frac{\Delta}{2} \left\{ \mathrm{e}^{2i\frac{g}{\omega}sin(\omega t)}|1\rangle\langle -1| + \mathrm{e}^{-2i\frac{g}{\omega}sin(\omega t)}|-1\rangle\langle 1| \right\} \psi_0 & \\ = -\frac{\Delta}{2} W \left\{ \mathrm{e}^{2i\frac{g}{\omega}sin(\omega t)}|0)(1| + \mathrm{e}^{-2i\frac{g}{\omega}sin(\omega t)}|1)(0| \right\} W\psi_0, & \end{aligned}$$

so we have only to solve the equation

$$i\frac{d}{dt}\phi = -\frac{\Delta}{2}\left\{\mathrm{e}^{2i\frac{g}{\omega}sin(\omega t)}\sigma_+ + \mathrm{e}^{-2i\frac{g}{\omega}sin(\omega t)}\sigma_-\right\}\phi, \quad \phi \equiv W\psi_0 \tag{8}$$

where $|0)(1| = \sigma_+$ and $|1)(0| = \sigma_-$.

Here we note that the solution ψ in (7) is given as follows

$$\psi(t) = W\left\{\mathrm{e}^{-i\frac{g}{\omega}sin(\omega t)}|0)(0| + \mathrm{e}^{i\frac{g}{\omega}sin(\omega t)}|1)(1|\right\}\phi(t) \tag{9}$$

with ϕ in (8).

To solve the equation (8) we set :

$$\phi = \begin{pmatrix} a \\ b \end{pmatrix}. \tag{10}$$

By substituting (10) into (8) we obtain the integral equations

$$a(t) = \alpha + i\frac{\Delta}{2}\int_0^t \mathrm{e}^{2i\frac{g}{\omega}sin(\omega s)}b(s)ds, \tag{11}$$

$$b(t) = \beta + i\frac{\Delta}{2}\int_0^t \mathrm{e}^{-2i\frac{g}{\omega}sin(\omega s)}a(s)ds, \tag{12}$$

or

$$a(t) = \alpha + i\frac{\Delta}{2}\beta\int_0^t \mathrm{e}^{2i\frac{g}{\omega}sin(\omega s)}ds - \frac{\Delta^2}{4}\int_0^t \mathrm{e}^{2i\frac{g}{\omega}sin(\omega s)}\left\{\int_0^s \mathrm{e}^{-2i\frac{g}{\omega}sin(\omega x)}a(x)dx\right\}ds, \tag{13}$$

$$b(t) = \beta + i\frac{\Delta}{2}\alpha\int_0^t \mathrm{e}^{-2i\frac{g}{\omega}sin(\omega s)}ds - \frac{\Delta^2}{4}\int_0^t \mathrm{e}^{-2i\frac{g}{\omega}sin(\omega s)}\left\{\int_0^s \mathrm{e}^{2i\frac{g}{\omega}sin(\omega x)}b(x)dx\right\}ds, \tag{14}$$

where α, β are integral constants. Therefore in the lowest order with respect to Δ, ϕ in (10) is given by

$$\phi = \begin{pmatrix} \alpha + i\frac{\Delta}{2}\beta\int_0^t \mathrm{e}^{2i\frac{g}{\omega}sin(\omega s)}ds \\ \beta + i\frac{\Delta}{2}\alpha\int_0^t \mathrm{e}^{-2i\frac{g}{\omega}sin(\omega s)}ds \end{pmatrix}.$$

From here we consider the physical condition. ψ in (9), therefore ϕ must be normalized ($\phi^\dagger\phi = 1$) in the lowest order, so we can set $|\alpha|^2 + |\beta|^2 = 1$. From (9) the approximate solutions that we are looking for are

$$\psi(t) = \frac{1}{\sqrt{2}}\begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}\begin{pmatrix} \mathrm{e}^{-i\frac{g}{\omega}sin(\omega t)}\left\{\alpha + i\frac{\Delta}{2}\beta\int_0^t \mathrm{e}^{2i\frac{g}{\omega}sin(\omega s)}ds\right\} \\ \mathrm{e}^{i\frac{g}{\omega}sin(\omega t)}\left\{\beta + i\frac{\Delta}{2}\alpha\int_0^t \mathrm{e}^{-2i\frac{g}{\omega}sin(\omega s)}ds\right\} \end{pmatrix}, \tag{15}$$

where $|\alpha|^2 + |\beta|^2 = 1$.

Finally we make a comment : If we write ψ above as

$$\psi = \begin{pmatrix} \psi_0 \\ \psi_1 \end{pmatrix},$$

then we have

$$|\psi_1|^2 = \frac{1}{2}\left(1 - \cos\left(\frac{2g}{\omega}\sin(\omega t)\right)\right) \tag{16}$$

under (B) if $\alpha = \beta = \mathrm{e}^{i\theta}/\sqrt{2}$.

Our solutions (15) are more or less well–known. However our (simple) method will become a starting point to find more complicated approximate solutions or to generalize the model from 2–levels to n–levels, [14].

The author would like to Marco Frasca for several useful comments.

References

[1] L. Allen and J. H. Eberly : Optical Resonance and Two–Level Atoms, Wiley, New York, 1975.

[2] A. Corney : Atomic and Laser Spectroscopy, Oxford University Press, 1977.

[3] M. S. Shahriar, P. Pradhan and J. Morzinski : Determination of the phase of an electromagnetic field via incoherent detection of fluorescence, quant-ph/0205120.

[4] M. S. Shahriar : Frequency Locking Via Phase Mapping of Remote Clocks Using Quantum Entanglement, quant-ph/0209064.

[5] M. S. Shahriar and P. Pradhan : Fundamental limitation on qubit operations due to the Bloch–Siegert Oscillation, in the Proceedings of the Quantum Communication, Measurement and Computing (QCMC'02), quant-ph/0212121.

[6] C. E. Creffield : Location of crossing in the spectrum of a driven two–level system, cond-mat/0301168.

[7] J. C. A. Barata and W. F. Wreszinski : Strong Coupling Theory of Two Level Atoms in Periodic Fields, Phys. Rev. Lett. 84(2000), 2112, physics/9906029.

[8] A. Santana, J. M. Gomez Llorente and V. Delgado : Semiclassical dressed states of two-level quantum systems driven by nonresonant and/or strong laser fields, J. Phys. B 34(2001), 2371, quant-ph/0011015.

[9] M. Frasca : Theory of dressed states in quantum optics, Phys. Rev. A 60(1999), 573, quant-ph/9811037.

[10] M. Frasca : Rabi oscillations and macroscopic quantum superposition states, Phys. Rev. A 66(2002), 023810, quant-ph/0111134.

[11] M. Frasca : A modern review of the two–level approximation, quant-ph/0209056.

[12] K. Fujii : Mathematical Structure of Rabi Oscillations in the Strong Coupling Regime, to appear in J. Phys. A, quant-ph/0203135.

[13] K. Fujii : N–Level System Interacting with Single Radiation Mode and Multi Cat States of Schrödinger in the Strong Coupling Regime, quant-ph/0210166.

[14] K. Fujii et al : N–Level System and Some Approximate Solutions in the Strong Coupling Regime (tentative), in progress.

[15] Y. Nakamura, Yu. A. Pashkin and J. S. Tsai : Rabi Oscillations in a Josephson–Junction Charge Two–Level System, Phys. Rev. Lett., 87(2001), 246601.